

## General Terms

- **Population Subgroups:**

Children:	Includes all children except for those who are patients with tuberculosis or transfusion recipients.
Homosexuals:	Includes people with known homosexual contacts such as "Men having sex with men."
Intravenous Drug Users/Needle Sharers:	Includes drug addicts.
Military/Armed Forces:	Includes police forces.
Other Non-Representative:	Includes those such as high-risk individuals and clients of sex workers, who do not fall into any of the other subpopulations.
Other Patients:	Includes all other patients except for pediatric patients, STI patients, TB patients, and Transfused patients.
Sex workers:	Includes other highly sexually active women such as bar workers.
Two Known Mixed Groups:	Includes those who fall into two different subpopulations but are reported as one. An example is pregnant women and sexually transmitted infected clinic patients who are reported as one group.

- **Data Type**

HIV	Human Immunodeficiency Virus (HIV) - non-specified probably HIV-1
HIV-1	HIV-1 (or HTLV III) - so specified
HIV-2	HIV-2 (or HTLV IV) - so specified
HIV-1 & 2	Co-infection of HIV-1 and HIV-2
HIV-1, 2	Overall combined of HIV-1 and HIV-2
AIDS	Acquired Immunodeficiency Syndrome

- **Test Type (These are based on an article from AIDS Volume 7, Number 1, January 1993 by Neil T. Constantine.**

ELISA	Enzyme-linked immunosorbent assay or EIA
ELISA*2, 3, ...	More than one ELISA test
IFA	Indirect immunofluorescent assay or IF
LIA	Line immunoassay
PCR	Polymerase chain reaction

RAPID	Rapid and simple assay
RAPID*2, 3, ...	More than one rapid assay
RIA	Radioimmuno assay
RIPA	Radioimmunoprecipitation assay
UNK	Unknown
WB	Western blot assay or IB

- **Specimen Type: Refers to what is collected and how it is collected from the donor.**

BW	Whole blood
BS	Blood spots
B	Blood (unspecified)
O	Oral/Saliva
U	Urine

- **Comments:** Common acronyms used in the “Comments for selected record” section

ANC	Antenatal clinic
AIS	AIDS Indicator Survey
ART	Antiretroviral therapy
AZT/ZDV	Zidovudine
BAIS	Botswana AIDS Impact Survey
BBS	Biological and Behavioral Survey
BED-CEIA	BED-capture enzyme immunoassay
CSW	Commercial sex worker
DHS	Demographic and Health Survey
DMPA	Depot medroxyprogesterone acetate
FBO	Faith-based organization
FSW	Female sex workers
FP	Family planning
FTC	Emtricitabine
GUD	Genital ulcer disease
HAART	Highly active antiretroviral therapy
HCT	HIV counseling and testing
HCV	Hepatitis C virus
HSV	Herpes simplex virus
IBBA	Integrated biological and behavioral assessment
IBBS	Integrated Bio-Behavioral Survey
MCH	Maternal and child health
MSM	Men having sex with men
NGO	Non-governmental organization
NVP	Nevirapine
PHIA	Population-Based HIV Impact Assessment
PMTCT	Prevention of mother-to-child transmission

PrEP	Pre-exposure prophylaxis
RDS	Respondent-driven sampling
SS	Sentinel Surveillance
STARHS	Serologic Testing Algorithm for Recent HIV Seroconversion
STI	Sexually transmitted infections
TDF	Tenofovir disoproxil fumarate
VCT	Voluntary counseling and testing
VMMC	Voluntary male medical circumcision